

FIPROtec[®] COMBO

50 mg/60 mg Spot-On Solution for Cats

READ THIS INFORMATION CAREFULLY BEFORE EACH USE. KEEP THIS LEAFLET IN THE OUTER CARTON FOR FUTURE REFERENCE.


WHAT IS FIPROtec[®] COMBO FOR?

FIPROtec[®] Combo spot-on solution for cats is used against infestations with fleas *Ctenocephalides* spp. (including their eggs and juvenile forms) and/or ticks (*Ixodes ricinus*, *Dermacentor variabilis*, *Rhipicephalus sanguineus*) and/or biting lice (*Felicola subrostratus*). Efficacy against fleas lasts for up to 4 weeks and against their multiplication for up to 6 weeks. Efficacy against ticks lasts for up to 2 weeks.

HOW DO I USE FIPROtec[®] COMBO?

Animals should be weighed accurately prior to treatment; do not use in kittens weighing less than 1 kg.

One tube should be given to each cat, applied to the skin as shown below.



1. Remove the pipette from its packaging. Hold the pipette in an upright position, twist and pull the cap off.
2. Turn the cap around and place the other end of the cap back on the pipette. Push and twist the cap to break the seal, then remove the cap from the pipette.

3. Apply to the skin at the base of the neck in front of the shoulder blades.
4. Part the coat until the skin is visible. Place the tip of the pipette onto the skin and squeeze the pipette several times to empty its contents completely and directly on the skin in one spot.

After treatment, you may notice transient clumping of hair or greasiness at the application site. It is important to make sure that the product is applied to an area where the animal cannot lick it off and to make sure that animals do not lick each other following treatment. If cats lick the product, it can cause them to salivate (drool) for a brief period of time.

In very rare cases, redness or discolouration of the skin, hair loss and itching may occur at the treatment site or on the cat in general. Excess salivation, reversible nervous signs (depression, hypersensitivity) or vomiting have been seen in exceptional cases. Bathing and/or shampooing of treated cats is not recommended. No information on the effect of bathing or the use of a shampoo on the effectiveness of the product in cats is available.

There may be an attachment of single ticks. For this reason, a transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.

WHAT IS IN FIPROtec[®] COMBO?

FIPROtec [®] COMBO CONTENTS 50 mg/60 mg spot-on solution cats			
Active substance		Other ingredients	
Amount of fipronil	50 mg	Butylhydroxyanisole (E320)	0.1 mg
Amount of S-methoprene	60 mg	Butylhydroxytoluene (E321)	0.05 mg

HOW OFTEN SHOULD I USE FIPROtec[®] COMBO?

Always leave a minimum period of 4 weeks between treatments. Adult fleas are killed for up to 4 weeks. Ticks are killed for up to 2 weeks. When a continued risk of flea and/or tick infestation is anticipated the treatment can be repeated every 4 weeks.

SHOULD I USE ANYTHING ELSE WITH FIPROtec[®] COMBO?

When treating infestations of parasites, all in-contact animals should be treated with an appropriate product at the same time. Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be

treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly.
Do not use simultaneously with other flea products which are applied directly on to the animal.

ANIMAL SAFETY WITH FIPROtec[®] COMBO

Do not use on kittens less than 8 weeks old or weighing less than 1 kg.

Do not use on rabbits as adverse reactions, including death, could occur.

Do not use on sick or convalescent animals.

Do not use if your cat has a hypersensitivity to fipronil, S-methoprene or the other ingredients.

Do not apply on wounds or damaged skin.

Avoid contact with your cat's eyes. If this occurs, immediately and thoroughly flush with water. If necessary, speak to your veterinary surgeon for advice.

In cats, the product can be used during pregnancy. No specific studies have been performed in lactating cats. Consult your veterinary surgeon before using the product in lactating cats or animals in contact with kittens less than 8 weeks old.

Speak to your veterinary surgeon if you notice any adverse effects following the use of FIPROtec[®] COMBO, even those not already listed in this package leaflet or you think that the medicine has not worked.

Speak to your veterinary surgeon if you notice any adverse effects following the use of FIPROtec[®] COMBO, even those not already listed in this package leaflet or you think that the medicine has not worked.

USER SAFETY WITH FIPROtec[®] COMBO

PLEASE READ THESE WARNINGS BEFORE EACH USE

Treated animals should not be handled or played with until the application site is dry. It is recommended that cats are treated during the early evening. Treated cats should not be allowed to sleep with owners, especially children.

People with a known hypersensitivity (allergy) to fipronil, S-methoprene or the other ingredients should avoid contact with the veterinary medicinal product.

This product can cause mucous membrane and eye irritation. Avoid getting the product on your skin, or in your mouth and eyes.

Wash off any product immediately with soap and water if skin contact occurs.

If the product gets in your eyes, immediately and thoroughly flush the eyes with water. If eye irritation persists seek medical advice and show the package leaflet or the label to the doctor.

If the product is accidentally swallowed, seek medical advice immediately and show the package leaflet to the doctor.

Do not smoke, drink or eat during application. Wash hands after use.

Keep pipettes in original packaging until ready to use.

GENERAL PRECAUTIONS FOR FIPROtec[®] COMBO

Keep out of the sight and reach of children.

For animal treatment only.

Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

HOW SHOULD FIPROtec[®] COMBO BE STORED?

Store below 30°C.

Store in the original package in order to protect from light and moisture.

HOW SHOULD FIPROtec[®] COMBO BE DISPOSED OF?

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Fipronil and S-methoprene may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty container.

Available in cartons containing 1,3 or 6 pipettes.

Not all pack sizes may be marketed.

424240

AVM-GSL Vm 01656/4122

Marketing authorisation holder and manufacturer for the batch release: KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

For further information on this product, please contact:
Distributor: Beaphar UK Ltd, Rook Tree Farm,
Withersfield Road, Great Wratting, Suffolk, CB9 7HD
Email: info@uk.beaphar.com
Tel: 0333 0066236.

Date on which the package leaflet was last approved: Maart 2021

13745-0

beaphar[®]

FIPROtec[®] COMBO

- Kills fleas, ticks & biting lice
- Prevents flea multiplication



424235

- - črna U
- - črna U (10%)
- - črna U (30%)
- - črna U (0%-100%)

Emb. mat.: NA.FIPROTEC COMBO RAZK CAT GB
Datum: 06.04.2018
Izdelaal: D. Kočjaž