

FIPROtec®

50mg Spot-On Solution for Cats
Fipronil

NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE:

Marketing authorisation holder:

Beaphar B.V., Drostenkamp 3, 8101 BX, Raalte, The Netherlands
T: +31 572 348 834, F: +31 572 348 835, E: hqb@beaphar.com

Manufacturer responsible for batch release:

Beaphar B.V., Oude Linderteseweg 9, 8102 EV Raalte, Netherlands
T: +31 572 348 834, F: +31 572 348 835, E: hqb@beaphar.com

ACTIVE SUBSTANCE AND OTHER INGREDIENTS:

1 pipette of 0.5 ml contains:

Active substance:

Fipronil 50.0 mg

Excipients:

Butylhydroxyanisole (E-320) 0.100 mg

Butylhydroxytoluene (E-321) 0.050 mg

Benzyl alcohol (E-1519) 142.500 mg

Clear colourless to slightly yellow solution.

TARGET SPECIES
Cats



WHAT DOES THIS PRODUCT DO?

BEAPHAR FIPROTEC® SPOT-ON FOR CATS kills fleas and ticks on your cat at the time of treatment, and continues to kill fleas for up to 5 weeks.

WHAT'S IN THIS PRODUCT?

Each pipette contains 50mg Fipronil (active ingredient), 0.1mg butylhydroxyanisole (E320), 0.05mg butylhydroxytoluene (E321) and 142.5mg benzyl alcohol (E1519).

IS THIS PRODUCT SUITABLE FOR MY CAT?

It is suitable for cats from 8 weeks of age, and weighing more than 1kg. Do not use if your cat is sick or is recovering from an illness. Do not use on pregnant or nursing cats. Ensure treated cats do not come into contact with kittens under 8 weeks of age. Do not use if your cat is allergic to any of the ingredients. For external use only.


Do not use on rabbits as serious reactions, including death, can result.

HOW DO I USE THIS PRODUCT?

Before every use, read the information on the inside of the box for full instructions and all warnings, including user warnings.


Use the easy-peel corners to remove a pipette from its blister.

Do not puncture the foil with scissors, knives or other sharp instruments, as this may damage the pipette inside.



Snip off the top of the pipette, part the fur between your cat's shoulder blades, place the tip of the pipette on the skin and apply the contents of one pipette to the skin surface, ideally split between two places about 2cm apart, one at the base of the head and a second 2-3cm further back. For continuous flea protection, re-apply the product once every 4 weeks, but not more often.

HOW TO APPLY FIPROtec® Spot-On:



HOW SHOULD I STORE AND DISPOSE OF THIS PRODUCT?

Keep the pipettes in the box, in order to protect from light, until ready to use them. Fipronil may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with product or empty pipettes. The product may have adverse effects on painted, varnished or other household surfaces or furnishings. Dispose of empty pipettes immediately after use in the household rubbish.

WHAT DO I NEED TO KNOW TO TREAT MY CAT SAFELY?

Do not use simultaneously with other flea products which are applied directly to the animal. Weigh your cat carefully before treating to ensure the product is safe to use. For external use only. Avoid contact with the animal's eyes. In case of accidental eye contact, immediately and thoroughly flush the eyes with water. Do not apply to wounds or damaged skin. Avoid excessive wetting of the fur. If stickiness occurs, this will disappear within 24 hours. Apply the product in a position where your cat cannot lick it off. Avoid frequent swimming or shampooing. In a multi-pet household, ensure treated animals cannot groom each other following treatment and that all animals are treated with a suitable insecticide. Fleas from pets often infest the animal's basket, bedding & regular resting areas, such as carpets and soft furnishings, which should be treated with a household insecticide and vacuumed regularly.

POSSIBLE ADVERSE REACTIONS:

If licked off, the liquid may cause a brief period of excessive drooling due to the taste of the solvent. Very rarely, temporary skin discolouration, hair loss or itching can occur at the site of treatment. In a few cases, skin sensitivity may be severe. Very rarely, a cat may temporarily appear depressed, vomit or show changes in its breathing. The risk of adverse reactions to the product may increase if overdosed. If you notice any effects after treatment, contact your veterinary surgeon.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- Common (more than 1 but less than 10 animals in 100 animals treated)
- Uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- Rare (more than 1 but less than 10 animals in 10,000 animals treated)
- Very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If a tick attaches to your cat, there is a possibility of a transmission of infectious diseases from the tick to your cat.

USER WARNINGS: WHAT PRECAUTIONS SHOULD I TAKE TO PROTECT MYSELF AND FAMILY?

For animal treatment only. Keep out of the sight and reach of children. Keep pipettes in this pack until you are ready to use. This product can cause mucous membrane and eye irritation. Therefore, contact of the product with mouth and eyes should be avoided. In case of accidental eye contact, rinse the eye carefully with plain water. If eye irritation continues, seek medical advice and show this pack to the doctor. If you have a known allergy to Fipronil or any of the other ingredients, avoid contact with this product. Avoid contents coming into contact with the fingers. If this occurs, wash your hands with soap & water. Do not smoke, drink or eat during application. Wash hands after use. Ingestion of the product is harmful. Prevent children getting access to the pipettes and discard the used pipettes immediately after applying the product. Children should not be allowed to play with treated animals, which should not be handled until after the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals should not be allowed to sleep with owners, especially children.

Date label text last approved: February 2022

OTHER INFORMATION

1, 2, 3, 4, 6 pipettes are packed in a cardboard box.
Not all pack sizes may be marketed.