



# Children's Dry Cough & Sore Throat Syrup

GLYCEROL  
& SUCROSE

FOR DAY & NIGHT USE



1 + year

### ⚠️ If you forget to give the medicine

If you forget a dose, give the next dose when needed. Do not give a double dose.

### Possible side-effects

This medicine is unlikely to cause side-effects unless your child is allergic to the ingredients. If your child experiences any of the following, stop giving the medicine and seek immediate medical help. See a doctor at once:

- Unexplained wheezing or difficulty in breathing, swelling of the face, lips, tongue, neck or throat (severe allergic reactions). If your child experiences any other side-effects not listed here or if you are not sure about anything, talk to your doctor or pharmacist.

### Reporting of side-effects

If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)  
By reporting side-effects you can help provide more information on the safety of this medicine.

### Storing this medicine

Do not give this medicine after the expiry date shown on the bottle. Keep the bottle tightly closed. There are no special storage instructions for this medicine. Keep out of the sight and reach of children. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. Retain carton for full instructions.

### Further information

What's in this medicine: The active ingredients in each 5 ml of medicine are: Glycerol 0.75 ml and Sucrose 1.7 g. Other ingredients are: Citric acid monohydrate, sodium benzoate (E211), anthocyanin, blackcurrant flavour, blackcurrant juice, liquid glucose and purified water.

What the medicine looks like: Benylin Children's Dry Cough & Sore Throat Syrup is a clear, purple liquid available in 125 ml bottles.

If you need more advice ask your pharmacist.

Marketing Authorisation Holder:  
McNeil Products Limited, Maidenhead,  
Berkshire, SL6 3UG, UK.  
Manufacturer: McNeil Iberica S.L.U.,  
28805 Madrid, Spain.

PL 15513/0392

This carton was revised in October 2015.  
Benylin is a registered trade mark.

Contains spoon



125 ml e



# Children's Dry Cough & Sore Throat Syrup

GLYCEROL  
& SUCROSE

FOR DAY & NIGHT USE



1 + year

### What this medicine is for

This medicine relieves irritating, dry coughs and sore throats.

### Before giving the medicine to your child

This product is suitable for most people, but a few people should not use it. If you are in any doubt contact your doctor or pharmacist.

### ❌ Do not give your child this medicine...

- If s/he is under 1 year old.
- If s/he is allergic to any of the ingredients.

If any of these bullet points apply, get advice from a doctor or pharmacist without using this medicine.

### ❌ Talk to your doctor or pharmacist...

- If your child has an intolerance to some sugars. If this applies to your child now or in the past, talk to a doctor or pharmacist.

### ⚠️ Some of the ingredients can cause problems

- This product contains 2.15 g glucose and 1.7 g sucrose in each 5 ml, which may be harmful to the teeth. If you have been told by your doctor that your child has an intolerance to some sugars, contact your doctor before using this medicine. This should be taken into account in patients with diabetes.
- Carbohydrate content of this medicine is 3.4 g per 5 ml.

### How to use this medicine

- Check the information below to see how much medicine to give.
- For oral use only. Do not give more medicine than the carton (and/or label) tells you to.
  - Do not overfill the spoon. There is a double-ended spoon in the pack. Check the cap seal is not broken before first use. If it is, do not take this medicine.

### 1 Children under 1 year

Do not give to children under 1 year old.

### 1 Children 1 year and over and Adults

Age	Dose
Children 1 - 5 years	One 5 ml spoonful.
Children 5 years and over and adults (including the elderly)	Two 5 ml spoonfuls.
<ul style="list-style-type: none"> <li>• Give the dose 3 or 4 times within 24 hours, if needed.</li> <li>• If your child does not feel better or starts to feel worse, tell your doctor or pharmacist.</li> </ul>	

- You can take this medicine if you are pregnant or breastfeeding.

### ⚠️ If anyone has taken too much

If anyone has taken too much of this product, contact a doctor or your nearest Accident & Emergency department (Casualty), taking this pack with you.