

PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**DISPRIN
Aspirin**

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to take it carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If symptoms persist or worsen consult your doctor.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this Leaflet:

1. What Disprin is and what it is used for
2. Before you take Disprin
3. How to take Disprin
4. Possible side effects
5. How to store Disprin
6. Further information

**1. What is Disprin and what it is used for**

The active ingredient which makes Disprin work is aspirin.

Aspirin provides relief by changing the body's response to pain, swelling and fever.

Disprin can be used for the relief of :

- | | | |
|---------------|---------------------------|-------------------------|
| * headache | *symptoms of cold and flu | *fever |
| * period pain | *sore throat | *sciatica and neuralgia |
| * migraine | *lumbago & rheumatic pain | *toothache |

2. Before you take Disprin

There is a possible association between aspirin and Reye's syndrome when given to children. Reye's syndrome is a very rare disease, which can be fatal. For this reason aspirin should not be given to children aged under 16 years, unless on the advice of a doctor.

Do not take Disprin if

- you suffer from haemophilia
- you have or have had a stomach ulcer
- you are under 16 (unless advised by your doctor).

Speak to your doctor or pharmacist before taking Disprin if

- you are allergic to aspirin or any of the other ingredients (see section 6)
- you suffer from asthma.

Can you take Disprin with other medicines?

The effect of treatment may be influenced if Disprin is taken together with medicines such as:

- anticoagulants (which thin the blood)
- uricosurics (which promote the elimination of uric acid, e.g. medicines for gout)
- medicine for high blood pressure
- medicine for organ rejection after transplantation
- medicine for cancer or rheumatoid arthritis
- ibuprofen (pain killer).

.3000107

Always speak to a pharmacist or doctor before taking aspirin with another medicine.

Pregnancy and breast feeding:

Ask your doctor for advice before taking any medicine.

Do not take Disprin if you are in the last 3 months of your pregnancy or if you are breastfeeding.

3. How to take Disprin

Dissolve the tablets in water before taking.

Adults, the elderly, children over 16: 2 to 3 tablets.

Repeat the dose every 4 hours if needed, **but do not exceed 13 tablets in 24 hours.**

Do not exceed the stated dose. If symptoms persist consult your doctor.

If you take more Disprin than you should:

If you start feeling sick or experience symptoms such as deafness, ringing in your ears, dizziness, abnormal sweating, warm hand and feet with racing pulse, an increased breathing rate, dehydration, you should seek immediate medical advice.

4. Possible side effects

After taking this product you may experience side effects such as:

- Signs of allergic reaction : asthma attack, breathing difficulties, skin rashes
- Signs of intestinal bleeding : vomiting blood or particles looking like coffee grounds, passing black tarry or bright red stools

If you experience any side effect, stop taking the medicine and seek medical help immediately.

Also you can help to make sure that medicines remain as safe as possible by reporting any unwanted side effects via the internet at www.mhra.gov.uk/yellowcard. Alternatively you can call Freephone 0808 100 3352 (available from 10 a.m. to 2 p.m. Mondays to Fridays) or fill in a paper form available from your local pharmacy.

5. How to store Disprin

- Store below 25°C in a dry place.
- Keep out of the reach and sight of children.
- Do not use after the expiry date indicated on the box, after EXP. This date refer to the last day of the month.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information**What Disprin contains:**

- **The active substance is aspirin.** Each dispersible tablet contains **300 mg of aspirin.** The other ingredients are calcium carbonate, maize starch, citric acid, talc, sodium lauryl sulphate, saccharin, crospovidone and lime flavour.
- Disprin tablets are white to off-white tablets. They are available in blister packs containing 16 or 32 tablets. The pack of 32 tablets is a pharmacy only medicine.

Marketing Authorisation Holder and manufacturer

Reckitt Benckiser Healthcare (UK) Ltd, Dansom Lane, Hull HU8 7DS.
PL 00063/0017

Last revised May 2013.

.3000107

UNIT HEIGHT = 170mm

UNIT HEIGHT = 170 mm

PRINT FREE AREA

PRINT FREE AREA

FRONT**BACK**

TR708697A
Dimensions:170x130mm

CUSTOMER INFO:
Minimum Point Size = 9.00pt