



How to use this medicine

Check the table below to see how much medicine to use.

Oraldene is for topical use only, which means that it should not be swallowed or taken internally.

i Children under 6 years old
Oraldene is not recommended for children under 6 years of age.

i Adults and Children aged 6 years and over

Age	Dose
Adults and children aged 6 years and over	Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day or as otherwise directed.

- Do not dilute.
- Do not swallow.
- Shake well before use.

i If anyone accidentally swallows this product

If anyone accidentally swallows this product, contact a doctor or your nearest Accident and Emergency department (Casualty) taking this pack with you.

i If you forget to use the medicine

Use the next dose as soon as you remember unless it is time to use it again. Do not use two quantities in one go.

This carton was revised August 2013. © Oraldene is a registered trade mark.


Product Licence holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.
Manufacturer: Famar Orléans, 5 avenue de Concy, 45071 Orléans, Cedex 2, France.



Oraldene®

Hexetidine

Medicated Relief for Mouth & Throat Infections



Possible side-effects

Oraldene can have side-effects, like all medicines, although these don't affect everyone and are usually mild. If you experience the following, stop using the medicine and talk to your doctor:

- Swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- Allergic reactions including itchy rash Irritation, tingling or numbness of the tongue and/or tissues inside the mouth Shortness of breath Cough Dry mouth Feeling sick (nausea) Sickness (vomiting)
- Salivary gland enlargement Taste disturbance or loss of taste Pain, swelling, redness or heat Tongue or tooth discolouration Blistering or ulceration

If you get any side effects, talk to your doctor, pharmacist or nurse.
Reporting of side-effects
If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side-effects you can help provide more information on the safety of this medicine.

Storing this medicine

Do not use if bottle seal is broken when purchased. Do not store above 30°C. Use within 6 months of opening. Protect from light. Keep in the outer container. Keep out of the reach and sight of children. Do not use your medicine after the date shown as an expiry date on the packaging. Retain carton for full instructions. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Further information

What's in this medicine?
The active ingredient is: Hexetidine 0.1 g/100 ml
Other ingredients are: Polysorbate 60, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122) sodium calcium edetate, sodium hydroxide (E524), and purified water.

What the medicine looks like
Oraldene is a clear red solution available in 100 ml and 200 ml bottles. It contains 0.1 g of the active ingredient in every 100 ml of medicine.

LOT
EXP
100 ml e
Medicated Relief for Mouth & Throat Infections
Hexetidine
Oraldene®

Oraldene®

Hexetidine

What the medicine is for

The mouthwash contains hexetidine, which is an antibacterial and antifungal and is used to help treat throat and mouth infections, including mouth ulcers, thrush, sore or bleeding gums (called gingivitis), bad breath (halitosis) or sore throat. It can also be used before and after dental surgery to help prevent infections.

Before using this medicine

This medicine is suitable for most people, but a few people should not use it. If you are in any doubt, talk to your doctor or pharmacist.

X Do not use this medicine...

- If you have ever had a bad reaction to Oraldene or any of its ingredients. If this applies to you, get advice from your doctor or pharmacist without using Oraldene.
- If your symptoms persist talk to your doctor or pharmacist.
- If you are pregnant or breast-feeding
 - Ask your doctor or pharmacist for advice before using Oraldene if you are pregnant or breast-feeding.
- Some of the ingredients can cause problems
 - Ethanol (96%) may cause irritation or dry skin if frequently applied to the skin.
 - Azorubin (E122) can cause allergic reactions, including asthma. Reactions are more common in people who are allergic to aspirin.
 - Sodium content: 0.016 g/100 ml.



BRaille

How to use this medicine
Check the table below to see how much medicine to use.
Oraldene is for topical use only, which means that it should not be swallowed or taken internally.

i Children under 6 years old
Oraldene is not recommended for children under 6 years of age.

i Adults and Children aged 6 years and over

Age	Dose
Adults and children aged 6 years and over	Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day or as otherwise directed.

- Do not dilute.
- Do not swallow.
- Shake well before use.

i If anyone accidentally swallows this product
If anyone accidentally swallows this product, contact a doctor or your nearest Accident and Emergency department (Casualty) taking this pack with you.

i If you forget to use the medicine
Use the next dose as soon as you remember unless it is time to use it again. Do not use two quantities in one go.

This carton was revised July 2013. © Oraldene is a registered trade mark.
Product Licence holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.
Manufacturer: Famar Orléans, 5 avenue de Concy, 45071 Orléans, Cedex 2, France.

McNeil Products Ltd
PL 15513/0067
MA 100/00401

oraldene

CONTENTS: The solution contains Hexetidine 0.1 g/100 ml. Other ingredients: Polysorbate 60, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122) sodium calcium edetate, sodium hydroxide (E524), and purified water.

Sodium content: 0.016 g/100 ml.

Oraldene has an antibacterial and antifungal action which relieves mouth and throat infections, including:

- **mouth ulcers**
- **sore and bleeding gums**
- **symptoms of sore throats**
- **bad breath**

Oraldene can also be used before and after dental surgery. Read and retain carton for full instructions.

Oraldene is a registered trade mark.

PL holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.

PL 15513/0067 MA 100/00401



Oraldene®

Hexetidine

Medicated Relief for Mouth & Throat Infections

Antibacterial Mouthwash to relieve
*gingivitis mouth ulcers
sore gums sore throat*
100 ml e

Do not use if bottle seal is broken when purchased. Keep out of the reach and sight of children.

DIRECTIONS: Mouthwash for topical use only. DO NOT SWALLOW. SHAKE WELL BEFORE USE.

ADULTS AND CHILDREN 6 YEARS AND OLDER

Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day. **CHILDREN UNDER 6 YEARS** - not recommended.

If symptoms persist discontinue treatment and consult your doctor or dentist.

- Do not dilute.
- Do not store above 30°C.

Use within 6 months of opening.

Protect from light. Keep container in the outer carton.

LOT

EXP



F-30841B-13-0004

F309419
13/0007

How to use this medicine

Check the table below to see how much medicine to use.

Oraldene is for topical use only, which means that it should not be swallowed or taken internally.

i Children under 6 years old

Oraldene is not recommended for children under 6 years of age.

i Adults and Children aged 6 years and over

Age	Dose
Adults and children aged 6 years and over	Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day or as otherwise directed.

- Do not dilute.
- Do not swallow.
- Shake well before use.

! If anyone accidentally swallows this product

If anyone accidentally swallows this product, contact a doctor or your nearest Accident and Emergency department (Casualty) taking this pack with you.

! If you forget to use the medicine

Use the next dose as soon as you remember unless it is time to use it again. Do not use two quantities in one go.

This carton was revised August 2013. © Oraldene is a registered trade mark.

Product Licence holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.
Manufacturer: Famar Orléans, 5 avenue de Concyr, 45071 Orleans, Cedex 2, France.




PL 15513/0067
MA 100/00401

Oraldene®

Hexetidine

Medicated Relief for Mouth & Throat Infections



Possible side-effects

Oraldene can have side-effects, like all medicines, although these don't affect everyone and are usually mild. If you experience the following, stop using the medicine and talk to your doctor:

- Swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- Allergic reactions including itchy rash
- Irritation, tingling or numbness of the tongue and/or tissues inside the mouth
- Shortness of breath
- Cough
- Dry mouth
- Feeling sick (nausea)
- Sickness (vomiting)
- Salivary gland enlargement
- Taste disturbance or loss of taste
- Pain, swelling, redness or heat
- Tongue or tooth discolouration
- Blistering or ulceration

If you get any side effects, talk to your doctor, pharmacist or nurse.

Reporting of side-effects

If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard.

By reporting side-effects you can help provide more information on the safety of this medicine.

Storing this medicine

Do not use if bottle seal is broken when purchased. Do not store above 30°C.

Use within 6 months of opening.

Protect from light.

Keep in the outer container.

Keep out of the reach and sight of children.

Do not use your medicine after the date shown as an expiry date on the packaging.

Retain carton for full instructions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.


Further information

What's in this medicine?

The active ingredient is: Hexetidine 0.1 g/100 ml
Other ingredients are: Polysorbate 60, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122) sodium calcium edetate, sodium hydroxide (E524), and purified water.

What the medicine looks like

Oraldene is a clear red solution available in 100 ml and 200 ml bottles. It contains 0.1 g of the active ingredient in every 100 ml of medicine.



Oraldene®

Hexetidine

What the medicine is for

The mouthwash contains hexetidine, which is an antibacterial and antifungal and is used to help treat throat and mouth infections, including mouth ulcers, thrush, sore or bleeding gums (called gingivitis), bad breath (halitosis) or sore throat. It can also be used before and after dental surgery to help prevent infections.

Before using this medicine

This medicine is suitable for most people, but a few people should not use it. If you are in any doubt, talk to your doctor or pharmacist.

X Do not use this medicine...


- If you have ever had a bad reaction to Oraldene or any of its ingredients. If this applies to you, get advice from your doctor or pharmacist without using Oraldene.
- If your symptoms persist talk to your doctor or pharmacist.

! If you are pregnant or breast-feeding

- Ask your doctor or pharmacist for advice before using Oraldene if you are pregnant or breast-feeding.

! Some of the ingredients can cause problems

- Ethanol (96%) may cause irritation or dry skin if frequently applied to the skin.
- Azorubin (E122) can cause allergic reactions, including asthma. Reactions are more common in people who are allergic to aspirin.
- Sodium content: 0.016 g/100 ml.



BRAILLE

How to use this medicine
Check the table below to see how much medicine to use.

Oraldene is for topical use only, which means that it should not be swallowed or taken internally.

i Children under 6 years old
Oraldene is not recommended for children under 6 years of age.

i Adults and Children aged 6 years and over

Age	Dose
Adults and children aged 6 years and over	Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day or as otherwise directed.

- Do not dilute.
- Do not swallow.
- Shake well before use.

! If anyone accidentally swallows this product
If anyone accidentally swallows this product, contact a doctor or your nearest Accident and Emergency department (Casualty) taking this pack with you.

! If you forget to use the medicine
Use the next dose as soon as you remember unless it is time to use it again. Do not use two quantities in one go.

This carton was revised July 2013. © Oraldene is a registered trade mark.

Product Licence holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.
Manufacturer: Famar Orléans, 5 avenue de Concyr, 45071 Orleans, Cedex 2, France.

PL 15513/0067
MA 100/00401

oraldene



CONTENTS: The solution contains Hexetidine 0.1g/100 ml. Other ingredients: Polysorbate 60, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122), sodium calcium edetate, sodium hydroxide (E524), and purified water.

Sodium content: 0.016 g/100 ml.

Oraldene has an antibacterial and antifungal action which relieves mouth and throat infections, including: **mouth ulcers • sore and bleeding gums • symptoms of sore throats • bad breath**. Oraldene can also be used before and after dental surgery.

Read and retain carton for full instructions.

Oraldene is a registered trade mark.

PL holder: McNeil Products Ltd, Maidenhead, Berkshire, SL6 3UG, UK.



PL 15513/0067
MA 100/00401

Oraldene®

Hexetidine

Medicated Relief for
Mouth & Throat
Infections

Antibacterial
Mouthwash to relieve
gingivitis mouth ulcers
sore gums sore throat
200 ml e

Do not use if bottle seal is broken when purchased.

Keep out of the reach and sight of children.

DIRECTIONS: Mouthwash for topical use only. DO NOT SWALLOW. SHAKE WELL BEFORE USE.

ADULTS AND CHILDREN 6 YEARS

AND OLDER

Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day. **CHILDREN UNDER 6 YEARS** - not recommended.

If symptoms persist discontinue treatment and consult your doctor or dentist.

- Do not dilute.
 - Do not store above 30°C.
- Use within 6 months of opening. Protect from light. Keep container in the outer carton.



FR0942013/0006

EXP

LOT