



Extra Advance 500 mg/65 mg Tablets

Paracetamol + Caffeine



Please read right through this leaflet before you start using this medicine.

This medicine is available without prescription, but you still need to use Panadol Extra Advance Tablets carefully to get the best results from them.

- Keep this leaflet, you may need to read it again.
- If you have any questions, or if there is anything you do not understand, ask your pharmacist.

In this leaflet:

1. What Panadol Extra Advance Tablets do
2. Check before you take Panadol Extra Advance Tablets
3. How to take Panadol Extra Advance Tablets
4. Possible side effects
5. How to store Panadol Extra Advance Tablets
6. Further information

1. What Panadol Extra Advance Tablets do
Panadol Extra Advance Tablets are faster absorbed than standard paracetamol and caffeine tablets. They are suitable for the relief of headache, migraine, backache, rheumatic pain, toothache and period pain. They also relieve the fever, aches and pains of colds and flu.

The tablets contain two active ingredients. Paracetamol is a painkiller and reduces your temperature when you have a fever. Caffeine acts to further help the effectiveness of paracetamol.



2. Check before you take Panadol Extra Advance Tablets



Do not take Panadol Extra Advance Tablets:

- if you have ever had an allergic reaction to paracetamol, caffeine or to any of the other ingredients (listed in Section 6)
- if you are taking other medicines containing paracetamol.



Ask your doctor before you take this medicine:

- if you have liver or kidney disease, including alcoholic liver disease.



If you are taking other medicines

Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly **metoclopramide** or **domperidone** (for **nausea** [feeling sick] or **vomiting** [being sick]) or **colestyramine** (to lower **blood cholesterol**). If you take **blood thinning drugs** (anticoagulants e.g. **warfarin**) and you need to take a pain reliever on a daily basis, **talk to your doctor** because of the risk of bleeding. But you can still take **occasional** doses of Panadol Extra Advance Tablets at the same time as anticoagulants.



Pregnancy and breast feeding

Due to the caffeine content of this product **it should not be used** if you are **pregnant** or **breast feeding**.



DFT-01192-3499567



3. How to take Panadol Extra Advance Tablets



Adults and children aged 16 years and over:

Swallow 2 tablets every 4 hours as needed. Do not take more than 8 tablets in 24 hours.

Children aged 12-15 years:

Swallow 1 tablet every 4 hours as needed. Do not give more than 4 tablets in 24 hours.



- **Do not take more frequently than every 4 hours.**
- **Do not take more than the recommended dose.**
- **Do not give to children under 12 years.**
- **Avoid too much caffeine in drinks like coffee and tea.**

High caffeine intake can cause **difficulty sleeping, shaking** and an **uncomfortable feeling in the chest.**

If you take too many tablets

Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

If your symptoms continue or your headache becomes persistent, **see your doctor.**

4. Possible side effects

Like all medicines, Panadol Extra Advance Tablets can have side effects but not everybody gets them. A small number of people have had side effects.

Stop taking the medicine and **tell your doctor immediately** if you experience:

- **Allergic reactions** which may be severe such as skin rash and itching sometimes with swelling of the mouth or face or shortness of breath
- **Skin rash or peeling, or mouth ulcers**
- **Breathing problems.** These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin
- Unexplained **bruising** or **bleeding**
- **Nausea**, sudden **weight loss, loss of appetite** and **yellowing of the eyes and skin.**



Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Panadol Extra Advance Tablets

Keep out of the sight and reach of children.

Do not use this medicine after the 'EXP' date shown on the pack. Store below 25°C.

6. Further information

Active ingredients: Each tablet contains Paracetamol 500 mg and Caffeine 65 mg.

Other ingredients: Starch pregelatinised, povidone k-25, calcium carbonate, crospovidone, alginate acid, magnesium stearate and sodium methyl (E 219), sodium ethyl (E 215) and sodium propyl (E 217) parahydroxybenzoates.

The tablet coating contains titanium dioxide (E 171), hypromellose, macrogol, polysorbate 80 and carnauba wax.

The parahydroxybenzoates may cause allergic reactions (possibly delayed).

Packs of Panadol Extra Advance Tablets contain 8, 14, or 32 tablets.

The marketing authorisation holder is GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

The manufacturer is GlaxoSmithKline Dungarvan Ltd., Co. Waterford, Ireland or Omega Pharma Manufacturing GmbH & Co. KG, Benzstraße 25, 71083, Herrenberg, Germany.

This leaflet was last revised in November 2018.

Trade marks are owned by or licensed to the GSK group of companies.

